



## Scientific Aspects of Agnihotra: Agriculture – Yield (part 10)

February 8, 2018

*By Dr. Ulrich Berk  
Germany*

In the last article we summarized four M.Sc. theses from Dharwad University, Karnataka, India, which give ample evidence that with Homa Organic Farming a healthy micro-flora and micro-fauna in soil is created, thus rejuvenating the soil. This is of utmost importance as nowadays conventional farming which uses a wide range of agrochemicals is destroying this micro-flora and micro-fauna, thus leaving behind a soil which is more and more exhausted, no longer a rich, living soil. Production comes down and even the supply of food for humanity is at stake, as reports from FAO, the [Food and Agriculture Organization of the United Nations](#) state.

A healthy soil is the capital for future generations to grow enough food; it is an investment in the future so to speak.

But equally important is production of enough food of good quality to feed the present population. How does Homa Organic Farming help with that? The four M.Sc. theses mentioned above also examined this question. Plants grown were soybeans, tomatoes, cabbage, and okra.

## Morphological Characteristics and Yield Parameters

The following table shows a significant increase in yield parameters.

We compare Homa Organic Farming (including seed treatment and application of Homa Biosol to plants and soil) with farming without Homa as control.

### Comparative Results: Homa vs. Non Homa

<b>Soybeans</b>		<b>Cabbage</b>	
	<b>Increase with Homa</b>		<b>Increase with Homa</b>
Total biomass per plant	28%	Plant height	40%
Straw yield	26%	No. of leaves per plant	23%
Grain yield	26%	Leaf area	13%
100 seeds weight	17%	Surface area of cabbage head	86%
Nodule count per plant	180%	Dry matter production	18%
Nodule dry weight per plant	172%	Yield per acre	15%

  

<b>Tomatoes</b>		<b>Okra</b>	
	<b>Increase with Homa</b>		<b>Increase with Homa</b>
No. of fruits per plant	33%	Plant height	39%
No. of leaves per plant	25%	No. of leaves per plant	60%
Dry matter	25%	Fruit weight	109%
Yield per plant	43%	Shelf life	125%
Yield per hectare	43%	Yield per hectare	28%

The table shows clearly that both morphological characteristics and yield improved considerably. These results you get when all methods of Homa Organic Farming are followed, including the treatment of seeds, foliar and soil application of Homa Biosol, and also spraying Agnihotra Ash water.

One of the results I want to mention especially: The development of root nodules in soybeans. Such nodules we find in leguminous plants like clovers, alfalfa, lupines, and peanuts as well as soybeans.

Nitrogen-fixing bacteria enter the roots of these legumes and multiply, thus forming root nodules.

In these nodules, nitrogen gas from the atmosphere (which is easily available there as 78% of our atmosphere is nitrogen) is fixed and made available to plants. When these plants die, the remaining nitrogen remains in the soil, thus improving soil quality and helping the growth of plants in the next season. This is of great importance, as nitrogen is often the limiting nutrient.

**How Homa Organic Farming (including the use of Homa Biosol) helps with the formation of root nodules becomes evident by the following photos.**



**Control without Homa**

**Nodules with Homa**

The four M.Sc. theses done at Dharwad University, Karnataka, India, also showed good effects of Homa Organic Farming on quality parameters of the foods grown and regarding reduction of pests and diseases. This will be discussed later.